
BD Insyte™, Insyte-W™, and Insyte-N™ IV Catheters

with BD Vialon™ Catheter Material

BD Insyte™ IV Catheter							
Ref#	Gauge	Length	Flowrate (mL/min)	Color			
381212	24 G (0.7 mm)	0.75 in. (19 mm)	25	Yellow			
381223	22 G (0.9 mm)	1.00 in. (25 mm)	35	Blue			
381233	20 G (1.1 mm)	1.00 in. (25 mm)	65	Pink			
381234	20 G (1.1 mm)	1.16 in. (30 mm)	60	Pink			
381237	20 G (1.1 mm)	1.88 in. (48 mm)	55	Pink			
381244	18 G (1.3 mm)	1.16 in. (30 mm)	105	Green			
381247	18 G (1.3 mm)	1.88 in. (48 mm)	95	Green			
381254	16 G (1.7 mm)	1.16 in. (30 mm)	220	Grey			
381257	16 G (1.7 mm)	1.77 in. (45 mm)	205	Grey			
381267	14 G (2.1 mm)	1.75 in. (45 mm)	330	Orange			



BD I nsyte- N™ IV Catheter						
Ref#	Gauge	Length	Flowrate (mL/min)	Color		
381211	24 G (0.7 mm)	0.56 in. (14 mm)	25	Yellow		
381311 (Winged)	24 G (0.7 mm)	0.56 in. (14 mm)	25	Yellow		

Shipping cases contain 200 units (4 boxes of 50)












bd.com


BD Insyte™ IV Catheter

Non-ported, straight IV catheter with BD Vialon™ Catheter Material



BD Insyte-W™ IV Catheter with Wings

All the benefits of BD Insyte™ IV Catheter with the **added benefits of wings**



Flexible wings are designed to stabilize the catheter

BD Insyte-N™ IV Catheter

Designed to accommodate **smaller veins**



Small gauge (24 G / 0.7 mm)

Short length (0.56 in. / 14 mm)

BD Instaflash™ Needle Technology

incorporates a notched needle, which is designed to confirm immediate vessel entry, improve first-stick success, and reduce painful hit-and-miss insertions


The BD Vialon™ Catheter Material Difference

More than 15 clinical studies conducted over 25 years have demonstrated that IV catheters made of BD Vialon™ Catheter Material demonstrated easier to insert or provide longer dwell times compared to FEP catheters.²⁻¹⁷

Improved patient comfort

Proprietary
BD Vialon™
Catheter Material
softens up to
70% in the vein.

BD Vialon™
Catheter Material
has greater kink
resistance than
FEP.²



"On average, Vialon remained in situ for 9 additional hours or 21% longer than Teflon before infiltrating."4

Reduced complications

BD Vialon™
Catheter Material
reduces the
chance of
mechanical
phlebitis by up to
50% compared to
FEP.³

References

- 1. Dougherty L, Lister SE. The Royal Marsden Hospital Manual of Clinical Nursing Procedures. 9th ed. Malden, MA: Wiley Blackwell Publishing; 2015.
- 2. Gaukroger PB, Roberts JG, Manners TA. Infusion thrombophlebitis: a prospective comparison of 645 Vialon and Teflon cannulae in anaesthetic and postoperative use. *Anaesthesia and Intensive Care*. 1988; 16:265-271.
- 3. Maki DG, Ringer M. Risk factors for infusion-related phlebitis with small peripheral venous catheters. *Annals of Internal Medicine*. 1991; 114: 845-854
- 4. Stanley MD, Meister E, Fuschuber K. Infiltration during intravenous therapy in neonates: comparison of Teflon® and Vialon® catheters. Southern Medical Journal. 1992; 85:883-886.
- 5. Jacquot C, Fauvage B, Bru JP, Croize J, Calop J. Effect of type of material on thrombophlebitis risk with peripheral venous catheters. *Ann Fr Anestha Réanim.* 1989; 8:3-7. [French language]
- 5. Fassolt A. Phlebitogenicity of venous catheters of Vialon. Infusionsther Klin Ernahr. 1985; 12(6): 282-2866.
- 7. Foster L, Wallis M, Paterson B, James H. A descriptive study of peripheral intravenous catheters in patients admitted to a pediatric unit in one Australian hospital. *J Infus Nurs*. 2002; 23(3):159-167.
- Karadag A, Gorgulu S. Effect of two different short peripheral catheter materials on phlebitis development. *J Infus Nurs*. 2003; 23(3):158-166.
- 9. Kerrison T, Woodhull J. Reducing the risk of thrombophlebitis. Comparison of Teflon and Vialon cannulae. *Professional Nurse*. 1994; 9(10): 662-666.
- 10. Lai KK. Safety of prolonging peripheral cannula and IV tubing use from 72 hours to 96 hours. AJIC. 1998;25(1):66-70.
- 11. McKee JM, Shell JA, Warren TA, Campbell VP. Complications of intravenous therapy: a randomized prospective study—Vialon vs. Teflon. *J Infus Nurs.* 1989;12(5): 288-295.
- 12. Myles PS, Buckland MR, Burnett WJ. Single versus double occlusive dressing technique to minimize infusion thrombophlebitis: Vialon® and Teflon® cannulae reassessed. *Anaesth Intens Care.* 1991; 19:525-529.
- 13. Stoddard GJ, Davis HT. Development of thrombophlebitis in peripheral veins with Vialon and PTFE-Teflon cannulas: a double-blind, randomised, controlled trial. *Ann R Coll Surg Engl.* 1993; 74:296-299.
- **14.** Shotkin JD, Lombardo F. Use of an indwelling peripheral catheter for 3-5 day chemotherapy administration in the outpatient setting. *J Infus Nurs.* 1996; 19(6): 315-320.
- 15. Stonehouse J, Butcher J. Phlebitis associated with peripheral cannulae. Professional Nurse. 1996; 12(1):51-54.
- 16. Szycher M. Biostability of polyurethane elastomers: a critical review. J Biomaterial Applications. 1998; 3: 297-402
- 17. Treuren BC, Galletly DC. A comparison of intravenous cannulae available in New Zealand. Anaesth Intens Care. 1990; 18(4): 540-546.